

# Phospho-p27 Kip 1 (Ser10) monoclonal antibody

Catalog:	MB11350
Cuturog.	111011000

Host: Ra

```
Rabbit
```

```
Reactivity: Human, Mouse, Rat
```

## **BackGround:**

The encoded protein binds to and prevents the activation of cyclin E-CDK2 or cyclin D-CDK4 complexes, and thus controls the cell cycle progression at G1. The degradation of this protein, which is triggered by its CDK dependent phosphorylation and subsequent ubiquitination by SCF complexes, is required for the cellular transition from quiescence to the proliferative state.

### **Product:**

50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA

Molecular Weight:

Calculated MW: 22 kDa; Observed MW: 27 kDa

**Swiss-Prot:** 

P46527

**Purification&Purity:** 

Affinity Purified

**Applications:** 

WB: 1/500-1/1000 IHC: 1/50-1/100 IP: 1/20

#### **Storage&Stability:**

Store at  $4 \,^{\circ}{\rm C}$  short term. Aliquot and store at  $-20 \,^{\circ}{\rm C}$  long term. Avoid freeze-thaw cycles.

#### **Isotype:**

IgG

**DATA:** 



Western blot analysis of Phospho-p27 KIP 1 in Hela, 3T3 lysates using Phospho-p27 Kip 1 antibody.



Western blot analysis of Phospho-p27 KIP 1 in C6 lysates using Phospho-p27 KIP 1 antibody. Immunohistochemistry analysis of paraffin-embedded Human breast

cancer using p27 KIP 1 antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.

#### Note:

For research use only, not for use in diagnostic procedure.

# Bioworld Technology, Inc.

 
 Add:
 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416,USA.

 Email:
 info@bioworlde.com

 Tel:
 6123263284

 Fax:
 6122933841

#### Bioworld technology, co. Ltd. Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China. Email: <u>info@biogot.com</u> Tel: 0086-025-68037686 Fax: 0086-025-68035151