

# Tec monoclonal antibody

Catalog: MB11685

Host: R

Rabbit

Reactivity:

# **BackGround:**

Non-receptor tyrosine kinase that contributes to signaling from many receptors and participates as a signal transducer in multiple downstream pathways, including regulation of the actin cytoskeleton. Plays a redundant role to ITK in regulation of the adaptive immune response. Regulates the development, function and differentiation of conventional T-cells and nonconventional NKT-cells. Required for TCR-dependent IL2 gene induction. Phosphorylates DOK1, one CD28-specific substrate, and contributes to CD28-signaling. Mediates signals that negatively regulate IL2RA expression induced by TCR cross-linking. Plays a redundant role to BTK in BCR-signaling for B-cell development and activation, especially by phosphorylating STAP1, a BCR-signaling protein. Required in mast cells for efficient cytokine production. Involved in both growth and differentiation mechanisms of myeloid cells through activation by the granulocyte colony-stimulating factor CSF3, a critical cytokine to promoting the growth, differentiation, and functional activation of myeloid cells. Participates in platelet signaling downstream of integrin activation. Cooperates with JAK2 through reciprocal phosphorylation to mediate cytokine-driven activation of FOS transcription. GRB10, a negative modifier of the FOS activation pathway, is another substrate of TEC. TEC is involved in G protein-coupled receptor- and integrin-mediated signalings in blood platelets. Plays a role in hepatocyte proliferation and liver regeneration and is involved in HGF-induced ERK signaling pathway. TEC regulates also FGF2 unconventional secretion (endoplasmic reticulum (ER)/Golgi-independent mechanism) under various physiological conditions through phosphorylation of FGF2 'Tyr-215'. May also be involved in the regulation of osteoclast differentiation.

#### **Product:**

50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA Molecular Weight: Calculated MW: 74 kDa; Observed MW: 58-75 kDa

Human

**Swiss-Prot:** 

P42680

**Purification&Purity:** 

Affinity Purified

**Applications:** 

WB: 1/500-1/1000 IF: 1/50-1/200

### **Storage&Stability:**

Store at  $4 \,^{\circ}{\rm C}$  short term. Aliquot and store at  $-20 \,^{\circ}{\rm C}$  long term. Avoid freeze-thaw cycles.

# **Isotype:**

# IgG

**DATA:** 



Western blot analysis of Tec in K562 lysates using Tec antibody.

Immunocytochemistry analysis of Tec in Hela using Tec antibody, and DAPI

#### Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.	
Add:	1660 South Highway 100, Suite 500 St. Louis Park,
	MN 55416,USA.
Email:	info@bioworlde.com
Tel:	6123263284
Fax:	6122933841

## Bioworld technology, co. Ltd. Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China. Email: <u>info@biogot.com</u> Tel: 0086-025-68037686 Fax: 0086-025-68035151