

CD86 Rat monoclonal antibody

| Catalog: MB65837 Host: Rat | Reactivity: Mouse |
|---|--|
| | |
| BackGround: | Swiss-Prot: |
| CD80 (B7-1, BB1) and CD86 (B7-2, B70) are member | s P42082 |
| of the B7 family of cell surface ligands that regulate | Purification&Purity: |
| cell activation and immune responses. CD80 is expressed | The monoclonal antibody was affinity-purified from rat |
| on activated antigen presenting cells, including dendritie | c antiserum by affinity-chromatography using |
| cells, B cells, monocytes, and macrophages. CD86 is ex | epitope-specific immunogen and the purity is $> 95\%$ (by |
| pressed on resting monocytes, dendritic cells, activated H | B SDS-PAGE). |
| lymphocytes, and can be further upregulated in the pres | Applications: |
| ence of inflammation. CD80 and CD86 are ligands for | r IF (1/50 - 1/200) |
| CD28, which functions as a T cell costimulatory receptor | Storage&Stability: |
| Interaction of CD28 with CD80 or CD86 provides the | Store at $4 ^{\circ}{\rm C}$ short term. Aliquot and store at $-20 ^{\circ}{\rm C}$ long |
| second signal required for na we T cell activation, T cel | term. Avoid neeze-maw cycles. |
| proliferation, and acquisition of effector functions. Alter | specificity. |
| natively, CD80 and CD86 also act as ligands to CTLA-4 | ' Recognizes mouse CD86 |
| which results in the downregulation of T cell activity. | DATA: |
| Product: | |

Note:

For research use only, not for use in diagnostic procedure.

dium azide.

Molecular Weight:

~ 34 kDa

Rat IgG2a kappa. Liquid in PBS, pH 7.3, and 0.02% so-