

TMPase monoclonal antibody

Catalog: MB66228

Host: Mouse

Reactivity: Human

BackGround:

Prostatic Acid Phosphatase (ACPP or PAP) is a member of the histidine acid phosphatase family. It is a non-specific phosphatase that is capable of dephosphorylating tyrosine residues as well as phospholipids under mildly acidic conditions. ACPP has ecto-5'-nucleotidase activity in pain-sensing neurons where it converts AMP to adenosine, suppressing the pain response (1,2). ACPP occurs as two isoforms that are both heavily glycosylated. The secreted phosphatase (sPAP) is found predominantly in the prostate and seminal plasma, while the cellular isoform (cPAP) is broadly expressed at very low levels and is associated with the plasma and lysosomal membranes (3-5). Cellular PAP has been shown to dephosphorylate ErbB2 at various tyrosine residues effectively terminating signaling. Furthermore, the physical interaction between cPAP and ErbB2 appears to regulate androgen sensitivity in prostate cancer cells. Loss of cPAP in androgen-sensitive prostate cancer cells results in the development of a castration-resistant phenotype suggesting that ACPP plays a significant role in prostate cancer cell growth. ACPP is expressed in metastatic cells arising from prostate cancer - especially in prostate-derived bone metastasis - suggesting that it may be a relevant diagnostic indicator of prostate cancer re-emergence in bone.

Product:

Mouse IgG1. Liquid in PBS containing 50% glycerol, 0.2% BSA and 0.01% sodium azide.

Molecular Weight:

~ 50 kDa

Swiss-Prot:

P15309

Purification&Purity:

The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

IHC (1/100 - 1/300)


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

Recognizes endogenous levels of TMPase protein.

DATA:



Immunohistochemical analysis of TMPase staining in human prostate formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151