

PRODUCT DATA SHEET

Bioworld Technology,Inc.

MLH1 Rabbit monoclonal antibody

Catalog: MB66267 Host: Rabbit Reactivity: Human, Mouse, Rat

BackGround:

Mismatch repair (MMR), a conserved process that involves correcting errors made during DNA synthesis, is crucial to the maintenance of genomic integrity. MLH1 is the human homologue of the E. coli MMR gene mutL. MMR requires recognition of a base mismatch or insertion/deletion loop by a MutS homolog followed by recruitment of a MutL heterodimeric complex consisting of MLH1 and PMS1 (MutL-γ), PMS2 (MutL-α), or MLH3 (MutL-γ). Other factors required for MMR in eukaryotes are EXO1, PCNA, RFC, RPA, DNA polymerases, and DNA ligase. Inactivation of the MLH1 gene causes genome instability and predisposition to cancer. The MLH1 gene is frequently mutated in hereditary nonpolyposis colon cancer (HNPCC). MLH1 also plays a role in meiotic recombination.

Product:

Liquid in 50mM Tris-Glycine (pH 7.4), 0.15M NaCl, 50% Glycerol, 0.01% Sodium azide and 0.05% BSA.

Molecular Weight:

~ 83 kDa

Swiss-Prot:

P40692

Purification&Purity:

The antibody was purified by immunogen affinity chromatography.

Applications:

WB (1/500 - 1/1000), IF/ICC (1/50 - 1/100), IP (1/10 - 1/50)


Storage&Stability:

Store at $4\,\mathrm{C}$ short term. Aliquot and store at -20 C long term. Avoid freeze-thaw cycles.


Specificity:

Recognizes endogenous levels of MLH1 protein.

DATA:



Western blot analysis of MLH1 expression in K562 (A), C6 (B), NIH3T3 (C), Hela (D) whole cell lysates.



Immunofluorescent analysis of MLH1 staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a hidified chamber. Cells were washed with PBST and incubated with a AF594-conjugated secondary antibody (red) in PBS at room temperature in the dark.

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park,

MN 55416,USA.

Email: <u>info@bioworlde.com</u>

Tel: 6123263284 Fax: 6122933841 Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046,

P. R. China.

Email: <u>info@biogot.com</u>
Tel: 0086-025-68037686
Fax: 0086-025-68035151