

CD74 monoclonal antibody

Catalog: MB66794

Host: Mouse

Reactivity: Human

BackGround:

The human histocompatibility leukocyte antigen (HLA) class II-associated invariant chain is composed of at least four polypeptides. One of these polypeptide chains is expressed as a membrane-bound subunit and has been designated CD74. The loading of peptide onto the class II MHC protein (MHC II) appears to be regulated by CD74, which associates with MHC II during its migration to the endosomal compartment, where class II binds peptide. CD74 is expressed by cells of both T lymphocyte and B lymphocyte lineages. In fact, CD74 is broadly expressed in normal B lymphocytes, regardless of their histocompatibility leukocyte antigen (HLA) phenotype, while a subset of peripheral T lymphocytes that are MHC II negative do not express CD74.

Product:

Mouse IgG2b kappa. Liquid in PBS, pH 7.3, 30% glycerol, and 0.01% sodium azide.

Molecular Weight:

~ 34 kDa

Swiss-Prot:

P04233

Purification&Purity:

This antibody is purified through a protein G column.

Applications:

WB (1/500 - 1/1000), FC (1/10 - 1/50)


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.


Specificity:

Recognizes endogenous levels of CD74 protein.

DATA:



Western blot analysis of CD74 expression in Daudi (A), Raji (B) whole cell lysates.



Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151