

PRODUCT DATA SHEET

Bioworld Biotech Co., Ltd

Tubulin α (9H4) monoclonal antibody

Catalog: MB8025 Host: Mouse Reactivity: Human, Mouse, Rat, Odontobutis obscurus

BackGround:

Tubulin is a major cytoskeleton component that has three distinct forms, designated α , β and γ Tubulin. α and β Tubulins form heterodimers, which multimerize to form a microtubule filament. γ Tubulin forms a soluble multiprotein particle with several other proteins. This particle, designated the gammasome, is required for nucleating microtubule filaments at the centrosome. In several organisms, numerous isoforms of the Tubulins exist that are encoded by different genes. The α and β isoforms undergo a variety of post-translational modifications, which may affect microtubule stability and protein interactions. High expression of class II β Tubulin has been seen in elongating axons, indicating a role in neurite outgrowth. Tubulins may also play a role in non-neuronal cell process formation

Product:

1 mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 55 KDa

Swiss-Prot:

O71U36/P68363

Purification&Purity:

The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:5000~1:20000 IF: 1:500~1000


Storage&Stability:

Store at $4\,\mathrm{C}$ short term. Aliquot and store at $-20\,\mathrm{C}$ long term. Avoid freeze-thaw cycles.

Specificity:

 α -tubulin (9H4) mAb detects endogenous levels of α -tubulin protein.

DATA:



Western blot (WB) analysis of α-tubulin (9H4) mAb at

1:5000/10000/20000 dilution


Lane1:The Kidney tissue lysate of Mouse(20ug)

Lane2:The Kidney tissue lysate of Rat(20ug)

Lane3:PC12 whole cell lysate(20ug)

Lane4:CT-26 whole cell lysate(20ug)

Lane5:HEK293T whole cell lysate(20ug)



Immunofluorescence analysis of RAW264.7 cells using α -tubulin (9H4) mAb at dilution of 1:1000 (40x lens).

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park,

MN 55416,USA.

Email: <u>info@bioworlde.com</u>

Tel: 6123263284 Fax: 6122933841 Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046,

P. R. China.

Email: <u>info@biogot.com</u>
Tel: 0086-025-68037686
Fax: 0086-025-68035151