

## BrdU monoclonal antibody

Catalog: MB9001

Host: Mouse

Reactivity: All

### BackGround:

The halogenated pyrimidine thymidine analog bromodeoxyuridine (BrdU) is incorporated into newly synthesized DNA strands of S-phase cells and is useful for estimating the fraction of cells in S-phase. Additionally, the analysis of the uptake of BrdU is a reliable method to quantitate the degree of DNA-synthesis. BrdU is also useful for studying sister chromatid exchange and to isolate nascent DNA. UV-induced excision-repair synthesis is one method for incorporating BrdU into cellular DNA. Anti-BrdU antibodies bind to the exposed BrdU in single-stranded DNA after a hydrochloric acid denaturation step or nuclease digestion. Protease antigen recovery is necessary for most tissues or cells fixed with crosslinking agents such as formalin but may decrease the specificity of BrdUrd immunodetection. The monoclonal antibody Bu20a against BrdU stains BrdU incorporated into the nuclei of a wide range of proliferating cell types including human tumors growing in nude mice and tonsil lymphoid.

### Product:

1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

### Molecular Weight:

N/A

### Swiss-Prot:

N/A

### Purification&Purity:

The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

### Applications:

ELISA: 0.015~ 0.05 µg/ml

Flow cytometry: 1~3 µg for 1x10<sup>6</sup> cells

ICC/IF: 1~3 µg/ml

### Storage&Stability:


The antibody is stable in lyophilized form if stored at -20 °C or below. The reconstituted antibody can be stored for 2-3 weeks at 2-8 °C. For long term storage, aliquot and store at -20 °C or below. Avoid repeated freezing and thawing cycles.

### Specificity:

The specificity of the antibody has been tested by competitive ELISA. Binding was inhibited by 5-bromo-2'-deoxyuridine (BrdU), 5-chloro-2'-deoxyuridine (CldU) and 5-iodo-2'-deoxyuridine (IdU). No cross-reactivity was observed with 5-fluoro-2'-deoxyuridine (FdU) and thymidine (T).

### DATA:

Immunocytochemistry/Immunofluorescence analysis of HeLa cells incorporated with BrdU using BrdU mAb, MB9001



### Note:

For research use only, not for use in diagnostic procedure.

### Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: [info@bioworld.com](mailto:info@bioworld.com)

Tel: 6123263284

Fax: 6122933841

### Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: [info@biogot.com](mailto:info@biogot.com)

Tel: 0086-025-68037686

Fax: 0086-025-68035151