

TfR mutant protein:W528A,recombinant(hFc Tag)

Cuturog. Ther	Catalog	;: N	CP012	24
---------------	---------	------	-------	----

Type: Human

293F

BackGround:

Cellular uptake of iron occurs via receptor-mediated endocytosis of ligand-occupied transferrin receptor into specialized endosomes . Endosomal acidification leads to iron release. The apotransferrin-receptor complex is then recycled to the cell surface with a return to neutral pH and the concomitant loss of affinity of apotransferrin for its receptor. Transferrin receptor is necessary for development of erythrocytes and the nervous system (By similarity). A second ligand, the heditary hemochromatosis protein HFE, competes for binding with transferrin for an overlapping C-terminal binding site. Positively regulates T and B cell proliferation through iron uptake. Acts as a lipid sensor that regulates mitochondrial fusion by regulating activation of the JNK pathway. When dietary levels of stearate (C18:0) are low, promotes activation of the JNK pathway, resulting in HUWE1-mediated ubiquitination and subsequent degradation of the mitofusin MFN2 and inhibition of mitochondrial fusion. When dietary levels of stearate (C18:0) are high, TFRC stearoylation inhibits activation of the JNK pathway and thus degradation of the mitofusin MFN2.

Product:	
PBS,pH7.4	
Size:	
100ug/1mg	
Swiss-Prot:	
P02786	
-	

Purification&Purity:

The protein was purified from 293F and the purity is > 95% (by SDS-PAGE).

Applications:

The protein has a calculated MW of 76 kDa.

Storage&Stability:


Source:

Store at $4 \,^{\circ}{\rm C}$ short term. Aliquot and store at $-20 \,^{\circ}{\rm C}$ long term. Avoid freeze-thaw cycles.


Protein length:

AA Leu 122 - Ile 550 (Accession # P02786)

DATA:



Line M: Protein Marker; Line 1: TfR-W528A (Trp mutation Ala) 2 μ g



Line M: Protein Marker; Line 1: TfR-W528A (Trp mutation Ala) Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add:	1660 South Highway 100, Suite 500 St. Louis Park,
	MN 55416,USA.
Email:	info@bioworlde.com
Tel:	6123263284
Fax:	6122933841

Bioworld technology, co. Ltd.

 Add:
 No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

 Email:
 info@biogot.com

 Tel:
 0086-025-68037686

 Fax:
 0086-025-68035151