

CD147 polyclonal antibody

Catalog: NCP0177P

Host: Rabbit

Reactivity: Human

BackGround:

Basigin (EMMPRIN, CD147) is a type I integral membrane receptor protein belonging to the immunoglobulin superfamily. Basigin is a glycosylated protein with four known isoforms, of which isoform 2 is the most abundantly expressed. Multiple functions have been ascribed to Basigin; foremost among these is stimulating the secretion of extracellular matrix metalloproteinases by adjacent fibroblasts, a function which has been implicated in promoting tumor progression. Research studies have shown that Basigin is overexpressed by many tumor cells, and its expression level may correlate with tumor malignancy. A recent study identified the BASIGIN gene as a regulatory target of Slug, suggesting a role for Basigin in the process of epithelial-mesenchymal transition. Basigin has also been identified as a marker for a subset of highly suppressive regulatory T cells, and as an obligate receptor for the malarial parasite *Plasmodium falciparum* on human erythrocytes.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2.

Molecular Weight:

~ 42 kDa

Swiss-Prot:

P35613

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:2000~1:5000

IP 1:50 - 1:200

IF 1:100 - 1:500

Storage&Stability:


Store at 4 °C short term. Aliquot and store at -20 °C long

term. Avoid freeze-thaw cycles.

Specificity:

CD147 polyclonal antibody detects endogenous levels of CD147 protein.


DATA:



Western blot (WB) analysis of CD147 polyclonal antibody at 1:5000 dilution

Lane1:A375 cell membrane lysate(10ug)


Lane2:SK-OVCAR3 cell membrane lysate(4ug)



Western blot (WB) analysis of CD147 polyclonal antibody at 1:2000 dilution

Lane1:EC9706 cell membrane lysate(25ug)

Lane2:A549 cell membrane lysate(25ug)



Immunofluores-

cence analysis of MG63 cells using CD147 pAb at dilution of 1:200 (40

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416,USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.


Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151

x lens).



1 is 30% input. The western blot was probed using CD147 pAb

#NCP0177P.

Note:

For research use only, not for use in diagnostic procedure.

Immunoprecipitation of SK-OVCAR3 cell lysates using CD147 pAb (Sepharose Bead Conjugate)#BD0048 (lane 2 and lane 3) and Nonspecific IgG Control (Sepharose Bead Conjugate)#BD0048 (lane 4) .Lane

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151