

CD50/ICAM-3 Recombinant Protein

Catalog: NCP0215

Host: E.coli

Tag: His-tag

BackGround:

Cell adhesion molecules (CAMs) are a family of closely related cell surface glycoproteins involved in cell-cell interactions during growth. These proteins are thought to play an important role in embryogenesis and development. ICAM-3, also designated CD50 and ICAM-R, is a type I membrane protein that is thought to regulate morphological changes during cell locomotion. ICAM-3 acts as a counter-receptor for the leukocyte Integrin $\alpha L/\beta 2$, and is known to activate T cells and polymorphonuclear leukocytes. ICAM-3 also binds to Moesin, via the cytoplasmic domain of ICAM-3. The expression of ICAM-3 is induced by RANTES, a chemoattractant known to activate T lymphocytes. ICAM-3 is also a major ligand for the leukocyte Integrin LFA-1 (CD11a/CD18).

Product:

PBS, 4M Urea, PH7.4

Molecular Weight:

~50kDa

Swiss-Prot:

P32942

Purification&Purity:

Transferred into competent cells and the supernatant was purified by NI column affinity chromatography and the purity is > 85% (by SDS-PAGE).

Restriction Sites:

NdeI-XhoI


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Expression Vector:

pet-22b(+)

DATA:



Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151