

CD112/NECTIN2 polyclonal antibody

Catalog: NCP0248P

Host: Rabbit

Reactivity: Human, Mouse

BackGround:

Modulator of T-cell signaling. Can be either a costimulator of T-cell function, or a coinhibitor, depending on the receptor it binds to. Upon binding to CD226, stimulates T-cell proliferation and cytokine production, including that of IL2, IL5, IL10, IL13, and IFNG. Upon interaction with PVRIG, inhibits T-cell proliferation. These interactions are competitive. Probable cell adhesion protein.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2.

Molecular Weight:

~ 60-70 kDa

Swiss-Prot:

Q92692

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:2000~1:5000

IF 1:50~1:200


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

CD112/NECTIN2 polyclonal antibody detects endogenous levels of CD112/NECTIN2 protein.

DATA:




Western blot (WB) analysis of CD112/NECTIN2 polyclonal antibody at 1:2000 dilution

Lane1:CT-26 whole cell lysate(30ug)

Lane2:H1792 whole cell lysate(20ug)

Lane3:HEK293T whole cell lysate(20ug)



Immunofluorescence analysis of HEK293T cells using CD112/NECTIN2 pAb at dilution of 1:200.

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151