

PRODUCT DATA SHEET



Bioworld Technology, Inc.

Recombinant Human IL-8 (77a.a.)

□rHuIL-8/CXCL8□

Catalog Number: PR1047

Source: Escherichia coli.

Quantity: 5µg/25µg/1.0mg

Description

Interleukin 8/CXCL8 was originally discovered and purified independently by a number of laboratories as a neutrophil chemotactic and activating factor. It was also referred to as neutrophil chemotactic factor (NCF), neutrophil activating protein (NAP), monocytederived neutrophil chemotactic factor (MDNCF), T-lymphocyte chemotactic factor (TCF), granulocyte chemotactic protein (GCP) and leukocyte adhesion inhibitor (LAI). Many cell types, including monocyte/macrophages, T cells, neutrophils, fibroblasts, endothelial cells, keratinocytes, hepatocytes, chondrocytes, and various tumor cell lines, can produce CXCL8 in response to a wide variety of pro-inflammatory stimuli such as exposure to IL-1, TNF, LPS, and viruses. CXCL8 is a member of the alpha (C-X-C) subfamily of chemokines, which also includes platelet factor 4, GRO, IP-10, etc.

Molecular Weight:

8,904 Da, a single, non-glycosylated polypeptide chain containing 77 amino acids.

Purity:

>95% by SDS-PAGE and HPLC analyses.

Biological Activity:

Fully biologically active when compared to standard. Determined by its ability to chemoattract human peripheral blood neutrophils using a concentration range of 25-150 ng/ml, corresponding to a Specific Activity of □6.7 x 10³ IU/mg.

Physical Appearance:

Sterile Filtered White lyophilized (freeze-dried) powder.

Formulation:

Lyophilized from a 0.2mm filtered concentrated solution in PBS, pH 7.4.

AA Sequence:

A V L P R S A K E L R C Q C I K T Y S K P F H P K F I K
E L R V I E S G P H C A N T E I I V K L S D G R E L C L
D P K E N W V Q R V V E K F L K R A E N S

Endotoxin:

Less than 1EU/mg of rHuIL-8 as determined by LAL method.

Reconstitution:

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1% BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at <-20°C. Further dilutions should be made in appropriate buffered solutions.

Storage:

This lyophilized preparation is stable at 2-8°C, but should be kept at -20°C for long term storage, preferably desiccated. Upon reconstitution, the preparation is stable for up to one week at 2-8°C. For maximal stability, apportion the reconstituted preparation into working aliquots and store at -20°C to -70°C. Avoid repeated freeze/thaw cycles.

Usage:

This material is offered by USA Bioworld biotech for research, laboratory or further evaluation purposes. NOT FOR HUMAN USE. Made in China

MADE IN CHINA

Bioworld Technology, Inc.

1660 South Highway 100, Suite 500 St. Louis Park, MN

55416, USA.

Tel: 6123263284

Email: info@bioworld.com

Fax: 6122933841

Bioworld technology, co, Ltd.

No 9, weidi road Qixia District Nanjing, 210046, P, R.China.

Tel: 0086-025-86371664

Email: info@biogot.com

Fax: 0086-025-86213570