PRODUCT DATA SHEET



Bioworld Technology, Inc.

Recombinant Human soluble CD40 Ligand (rHusCD40L)

Catalog Number: PR1113 Source: Escherichia coli. Quantity: 10µg/50µg/1.0mg

Description

CD40 ligand, CD40L (also known as CD154, TRAP or gp39), is a 261 amino acid type II transmembrane glycoprotein belonging to the TNF family, CD40L is expressed predominantly on activated CD4+ T lymphocytes, and also found in other types of cells, like NK cells, mast cells, basophils and eosinophils. Human CD40L shares 78% amino acid identity with its murine counterpart. The receptor of CD40L is CD40, a type I transmembrane glycoprotein belonging to the TNF receptor family. CD40 is expressed on B lymphocytes, monocytes, dendritic cells and thymic epithelium. Although all monomeric, dimeric and trimeric forms of soluble CD40L can bind to CD40, the trimeric form of soluble CD40L has the most potent biological activity through oligomerization of cell surface CD40, a common feature of TNF receptor family members. CD40L mediates a range of activities on B cells including induction of activation-associated surface antigen, entry into cell cycle, isotype switching and Ig secretion and memory generation. CD40-CD40L interaction also plays important roles in monocyte activation and dendritic cell maturation.

Molecular Weight:

Approximately 16.3 kDa. a single non-glycosylated polypeptide chain containing 149 amino acids.

Purity:

>95% by SDS-PAGE and HPLC analyses.

Biological Activity:

Fully biologically active when compared to standard. The ED50 as determined by the dose-dependant stimulation of IL-12 IL-8 induction by PMB (Peripheral Mononuclear) cells

was found to be □10 ng/ml, corresponding to a specific activity of $\Box 1 \times 106 \text{ IU/mg}$.

Physical Appearance:

Sterile Filtered White lyophilized (freeze-dried) powder.

Formulation:

Lyophilized from a 0.2mm filtered concentrated solution in PBS, pH 7.0.

AA Sequence:

M Q K G D Q N P Q I A A H V I S E A S S K T T S V L Q WAEKGYYTMSNNLVTLENGKQLTVKR QGLYYIYAQVTFCSNREASSQAPFIASL WLKSPGRFERILLRAANTHSSAKPCGQ Q S I H L G G V F E L Q P G A S V F V N V T D P S Q V SHGTGFTSFGLLKL

Endotoxin:

Less than 1EU/mg of rHusCD40L as determined by LAL method.

Reconstitution:

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1% BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at <-20°C. Further dilutions should be made in appropriate buffered solutions.

Storage:

Tel: 0086-025-86371664

This lyophilized preparation is stable at 2-8°C, but should be kept at -20°C for long term storage, preferably desiccated. Upon reconstitution, the preparation is stable for up to one week at 2-8°C. For maximal stability, apportion the reconsti-

MADE IN CHINA

Bioworld technology, co, Ltd.

No 9, weidi road Qixia District Nanjing, 210046, P, R.China.

Email: info@biogot.com Fax:0086-025-86213570

Bioworld Technology, Inc.

Tel: 6123263284

Fax: 6122933841

PRODUCT DATA SHEET

tuted preparation into working aliquots and store at -20°C to -70°C. Avoid repeated freeze/thaw cycles.

Usage:

This material is offered by USA Bioworld biotech for research, laboratory or further evaluation purposes. NOT FOR HUMAN USE. Made in China

1660 South Highway 100, Suite 500 St. Louis Park,MN 55416,USA. Email: info@bioworlde.com

Tel: 6123263284 Fax: 6122933841

MADE IN CHINA

Bioworld technology, co, Ltd.

Tel: 0086-025-86371664

No 9, weidi road Qixia District Nanjing, 210046, P, R.China.

Email: info@biogot.com Fax:0086-025-86213570