

PRODUCT DATA SHEET



Bioworld Technology, Inc.

Recombinant Mouse Leukemia inhibitory factor(rMuLIF)

Catalog Number: PR2005

Source: Escherichia coli.

Quantity: 5µg/10µg/1.0mg

Description

Leukemia Inhibitory Factor (LIF) is a lymphoid factor which promotes long-term maintenance of embryonic stem cells by suppressing spontaneous differentiation. LIF has a number of other activities including cholinergic neuron differentiation, control of stem cell pluripotency, bone and fat metabolism, mitogenesis of certain factor dependent cell lines and promotion of megakaryocyte production in vivo. Mouse LIF is a 20 kDa protein containing 181 amino acid residues.

Molecular Weight:

Approximately 20 kDa, a single non-glycosylated polypeptide chain containing 181 amino acids.

Purity:

>98% by SDS-PAGE and HPLC analyses.

Biological Activity:

The activity of mouse LIF is determined by the ability to induce differentiation of murine M1 myeloid leukemic cells. The minimum detectable concentration of mouse LIF in this assay is 0.5 ng/mL. The specific activity is >1 x 10⁸ units/mg, where 50 units is defined as the amount of mouse LIF required to induce differentiation in 50% of the M1 colonies in 1 mL agar cultures.

Physical Appearance:

Sterile Filtered White lyophilized (freeze-dried) powder.

Formulation:

Lyophilized from a 0.2µm filtered concentrated solution in 20mM PB, pH 7.4, with 0.02% TWEEN 20.

AA Sequence:

M S P L P I T P V N A T C A I R H P C H G N L M N Q I
K N Q L A Q L N G S A N A L F I S Y Y T A Q G E P F P
N N V E K L C A P N M T D F P S F H G N G T E K T K L
V E L Y R M V A Y L S A S L T N I T R D Q K V L N P T
A V S L Q V K L N A T I D V M R G L L S N V L C R L C
N K Y R V G H V D V P P V P D H S D K E A F Q R K K
L G C Q L L G T Y K Q V I S V V V Q A F

Endotoxin:

Less than 1EU/mg of rmLIF as determined by LAL method.

Reconstitution:

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1% BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at <-20°C. Further dilutions should be made in appropriate buffered solutions.

Storage:

This lyophilized preparation is stable at 2-8°C, but should be kept at -20°C for long term storage, preferably desiccated. Upon reconstitution, the preparation is stable for up to one week at 2-8°C. For maximal stability, apportion the reconstituted preparation into working aliquots and store at -20°C to -70°C. Avoid repeated freeze/thaw cycles.

Usage:

This material is offered by USA Bioworld biotech for research, laboratory or further evaluation purposes. NOT FOR HUMAN USE. Made in China

MADE IN CHINA

Bioworld Technology, Inc.

1660 South Highway 100, Suite 500 St. Louis Park, MN

55416, USA.

Tel: 6123263284

Email: info@bioworld.com

Fax: 6122933841

Bioworld technology, co, Ltd.

No 9, weidi road Qixia District Nanjing, 210046, P, R.China.

Tel: 0086-025-86371664

Email: info@biogot.com

Fax: 0086-025-86213570