

Bioworld Technology, Inc.

Recombinant Murine CXCL16 (rMuCXCL16)

Catalog Number: PR2008

Source: Escherichia coli.

Quantity:5µg/25µg/1.0mg

Description

Mouse CXCL16 (CXC chemokine 16) is a nonELR motifcontaining CXC chemokine with a transmembrane domain. CX3CL1/Fractalkine and CXCL16 are the only two transmembrane chemokines within the superfamily. Mouse CXCL16 cDNA encodes a 246 amino acid (aa) precursor protein with a putative 26 aa residue signal peptide, an 88 aa residue chemokine domain, an 87 aa residue mucinlike spacer region, a 22 aa residue transmembrane domain, and a 23 aa residue cytoplasmic tail. Mouse and human CXCL16 share 49% overall aa identity and 70% similarity in the chemokine domains. Mouse CXCL16 is produced by dendritic cells in lymphoid organ T cell zones and by cells in the splenic red pulp both as membranebound and soluble forms. Based on northern blot analysis,CXCL16 is also expressed in some nonlymphoid tissues such as lung, small intestine and kidney. The receptor for CXCL16 has been identified as CXCR6/Bonzo (STRL33 and TYMSTR), a receptor previously shown to be a coreceptor for HIV entry.CXCR6 is expressed on naive CD8 cells, natural

killer T cells and activated CD8 and CD4 T cells.

Molecular Weight:

Recombinant murine CXCL16 is a 9.9 kDa protein containing 88 amino acid residues.

Purity:

>98% by SDS-PAGE and HPLC analyses.

Biological Activity:

Fully biologically active when compared to standard. Determined by its ability to chemoattract murine lymphocytes using a concentration of 20-1000 ng/ml, corresponding to a Specific Activity of \Box 100 IU/mg.

Physical Appearance:

Sterile Filtered White lyophilized (freeze-dried) powder.

Formulation:

Lyophilized from a 0.2mm filtered concentrated solution in PBS.

AA Sequence:

NQGSVAGSCSCDRTISSGTQIPQGTLDH IRKYLKAFHRCPFFIRFQLQSKSVCGGS QDQWVRELVDCFERKECGTGHGKSFH HQKHLP

Endotoxin:

Less than 1EU/mg of rMuCXCL16 as determined by LAL method.

Reconstitution:

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1% BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at <-20°C. Further dilutions should be made in appropriate buffered solutions.

Storage:

This lyophilized preparation is stable at 2-8°C, but should be kept at -20°C for long term storage, preferably desiccated. Upon reconstitution, the preparation is stable for up to one

Bioworld Technology, Inc.

1660 South Highway 100, Suite 500 St. Louis Park,MN55416,USA.Email: info@bioworlde.comTel: 6123263284Fax: 6122933841

MADE IN CHINA Bioworld technology, co, Ltd. No 9, weidi road Qixia District Nanjing, 210046, P, R.China. Email: info@biogot.com Tel: 0086-025-86371664 Fax:0086-025-86213570 week at 2-8°C. For maximal stability, apportion the reconstituted preparation into working aliquots and store at -20°C to -70°C. Avoid repeated freeze/thaw cycles. Usage:

This material is offered by USA Bioworld biotech for research, laboratory or further evaluation purposes. NOT FOR HUMAN USE. Made in China

Bioworld Technology, Inc. h Highway 100, Suite 500 St. Louis Park.

 1660 South Highway 100, Suite 500 St. Louis Park,MN

 55416,USA.
 Email: info@bioworlde.com

 Tel: 6123263284
 Fax: 6122933841